



Robustness of a cross contamination model describing transfer of pathogens during grinding of meat

Møller, Cleide Oliveira de Almeida; Sant'Ana, A. S.; Hansen, Solvej Katrine Holm; Nauta, Maarten; Silva, L. P.; Alvarenga, V. O.; Maffei, D.; Pacheco, F.; Lopes, J.; Franco, B. D. G. M.

Total number of authors:

12

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Møller, C. O. D. A. (Author), Sant'Ana, A. S. (Author), Hansen, S. K. H. (Author), Nauta, M. (Author), Silva, L. P. (Author), Alvarenga, V. O. (Author), Maffei, D. (Author), Pacheco, F. (Author), Lopes, J. (Author), Franco, B. D. G. M. (Author), Aabo, S. (Author), & Hansen, T. B. (Author). (2015). Robustness of a cross contamination model describing transfer of pathogens during grinding of meat. Sound/Visual production (digital), Technical University of Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Robustness of a cross contamination model describing transfer of pathogens during grinding of meat

Møller C.O. de A.^a, Sant'Ana A.S.^{b,c}, Hansen S.K.H.^a, Nauta M.J.^a, Silva L.P.^c,
Alvarenga V.O.^{b,c}, Maffei D.^b, Pacheco F.^b, Lopes J.^b, Franco B.D.G.M.^b, Aabo S.
^a, Hansen T.B.^a

^aTechnical University of Denmark, Mørkhøj Bygade 19, Søborg, DK 2860, Denmark

^bUniversity of São Paulo, Avenida Professor Lineu Prestes 580, São Paulo, SP 05508-000, Brazil

^cState University of Campinas, Rua Monteiro Lobato 80, Campinas, SP 13083-862, Brazil

Correspondence

Cleide Møller. E-mail: clemo@food.dtu.dk



ORIGINAL ARTICLE

Modelling transfer of *Salmonella* Typhimurium DT104 during simulation of grinding of pork

C.O.A. Møller¹, M.J. Nauta¹, B.B. Christensen², P. Dalgaard³ and T.B. Hansen¹

¹ National Food Institute, Technical University of Denmark, Søborg, Denmark

² Department of Food Science, Faculty of Life Sciences, University of Copenhagen, Frederiksberg C, Copenhagen, Denmark

³ National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark

Journal of Applied Microbiology

Volume 112, Issue 1, January 2012, Pages: 90–98

Why Robustness?

Can the model predict /describe transfer of different pathogens during grinding of different food matrices using different equipment at different temperatures?

Aim


To investigate the robustness of a transfer model by three different criteria

1) ASZ

2) QMRA approach

3) TTP%

17 + 3 grinding trials





Different grinding conditions

- Type of grinder
 - 2 different degrees of roughness in the surfaces
- Type of meat matrice
 - Pork
 - Beef
- Size of pieces of meat
 - 50 g
 - 157 – 324 g
- Number of pieces of meat
 - 10 - 100
- Pathogen
 - Single strain of *S. Enteritidis* 54
 - Cocktail of *S. Enteritidis*
 - *S. Typhimurium* DT104
 - *L. monocytogenes*
- Room temperature
 - 4 °C
 - 19-27 °C
- Sharpness of grinder knife

Robustness assessed by Acceptable simulation Zone

- Applying the parameters suggested by Møller et al. in 2012 to simulate the cross contamination events observed in the remaining nineteen trials.
- In this study, ASZ of $\pm 0.5 \log_{10}\text{CFU}$ per portion.

70 % of observations inside the ASZ



Robustness assessed by a QMRA approach

International Journal of Food Microbiology

Volume 196, Issue 1, 2015, Pages: 109–125



Contents lists available at [ScienceDirect](#)

International Journal of Food Microbiology

journal homepage: www.elsevier.com/locate/ijfoodmicro



Risk assessment of *Salmonella* in Danish meatballs produced in the catering sector




Cleide O. de A. Møller^a, Maarten J. Nauta^a, Donald W. Schaffner^b, Paw Dalgaard^c,
Bjarke B. Christensen^{a,1}, Tina B. Hansen^{a,*}

^a National Food Institute, Technical University of Denmark, Mørkhøj Bygade 19, DK-2860 Søborg, Denmark

^b Department of Food Science, School of Environmental and Biological Science, Rutgers University, 65 Dudley Road, Food Science Building, Room 207, New Brunswick, NJ 08901-8520, USA


^c National Food Institute, Technical University of Denmark, Søtofts Plads, Bygning 221, DK-2800 Kgs. Lyngby, Denmark

Relative risk from different grinding parameters comparing to the baseline of Møller et al 2012.



Robustness assessed by Total Transfer Potential (TTP %)

$$TTP = \frac{\sum_{i=1}^{\infty} M_i}{S_1} = \left(1 - a_2 \left(1 - a_1 - \frac{b_2}{c_3 + b_2(1 - c_3)} \right) \right) \times 100\%$$



Correlation between all three approaches applied for model evaluation

> 1.1 log₁₀CFU > 2.1 x 10⁻³ > 60 %

Table 2. Summary of performance results (with extreme values in **bold**) obtained with three different approaches for evaluating performance of the model proposed by Møller et al. (2012)¹ describing the transfer of pathogens during meat grinding.

Trial	Parameter estimates from fitting					RMSE ^b	Size of ASZ ^c to include 100 % of the predictions (± CFU/portion of meat)	Absolute Risk ^e Estimates x 10 ⁻³	TTP % ^f
	a ₁	b ₁	a ₂	b ₂	1-c ₃				
1	0.0558	0.0544	0.4946	0.4175	1.0000	1.2038	2.0	3.02	103
2	0.0503	0.0454	0.1707	0.3522	1.0000	1.1536	2.0	2.87	101
3	0.0335	0.0569	0.0612	0.1485	1.0000	1.0754	2.0	2.94	100
4	0.0041	0.0471	0.8778	0.0570	0.4641	1.1018	0.8	1.01	21
5	0.0118	0.0652	0.7787	0.0927	0.5058	1.0760	1.4	1.56	36
6	0.0104	0.0677	0.8330	0.0588	0.3004	1.1079	1.2	1.10	24
7	0.0092	0.1472	0.8697	0.0261	0.1459	1.1879	1.1	0.86	16
8	0.0107	0.0265	0.9388	0.0253	0.4024	1.1508	1.2	0.62	11
9	0.0167	0.0061	0.5386	0.0579	0.8083	1.2202	2.0	2.21	60
10	0.4227	0.0052	0.4250	0.2715	0.5905	2.0930	0.7	1.94	96
11	0.0047	0.0358	0.6549	0.1251	0.5479	1.1744	1.1	1.92	51
12	0.0799	0.3165	0.2611	1.0000	0.0000	1.4371	1.4	3.04	102
13	0.0293	0.2433	0.8785	0.1648	0.3284	1.1196	1.2	1.44	35
14	0.0054	0.1127	0.5721	0.1653	0.6541	1.1802	1.5	2.22	64
15	0.0054	0.1584	0.7369	0.0484	0.6276	1.1909	1.1	1.54	36
16	0.0130	0.1134	0.5289	0.1304	0.4963	1.1451	1.1	2.24	60
17	0.0142	0.1022	0.7746	0.1767	0.1588	1.2773	2.2	1.59	39
18 ^a	0.0008	0.0655	0.7924	0.1331	0.2475	1.2029	0.6	1.41	34
19 ^a	0.0020	0.0809	0.8166	0.0555	0.3692	1.1345	0.6	1.14	25
20 ^a	0.0010	0.0275	0.8909	0.0558	0.4887	1.1378	NA ^d	0.96	20

^a data published by Møller et al. (2012).

^b Root Mean Sum of squared Errors.

^c ASZ – Acceptable Simulation Zone, proposed by Oscar in 2005² and tested by Møller et al. in 2013³.

^d NA – Not Applicable, because the parameters applied to access the ASZ were obtained by the fitting of trial 20.

^e risk estimates from scenarios testing different sets of transfer parameters (Table 2), and using the QMRA of *Salmonella* in meatball processing model (Møller et al., 2015⁴) at low concentration and prevalence of the pathogen.

^f Calculated with the equation derived from Møller et al. (2012)¹. It indicates the percentage (%) of CFU of *Salmonella*, from the contaminated grinded pieces that ends up in the total minced meat, assuming that the grinding process will continue forever.

Robustness of a cross contamination model describing transfer of pathogens during meat grinding

Final Considerations

- Applying the **Acceptable Simulation Zone** the cross contamination parameters suggested by Møller et al. (2012) **are not** able to describe all nineteen observed trials.
- The observed values had its cross contamination event **well described when fitted** to the model structure proposed by Møller et al. (2012).
- Parameter estimates obtained by fitting observed trials performed at different conditions, such as size of processed pieces of meat and number of processed slices, **may not be applied** to describe cross contamination of **unlike processing**.
- The **QMRA**, and the **Total Transfer Potential**, revealed that the risk may be reduced when the grinding is performed in a grinder made of stainless steel, using well-sharpened knives, and at room temperatures lower than 4°C.

Acknowledgements




Skilful technical assistance was provided by the colleagues from:

Division of Microbiology and Production at the National Food
Institute of the **Technical University of Denmark**

Acknowledgements

Skilful technical assistance was provided by the colleagues from the:

Department of Food and Experimental Nutrition,
Faculty of Pharmaceutical Sciences, **University of São Paulo**



Quantitative Food Microbiology Laboratory (LMQA)
Department of Food Science in the Faculty of Food Engineering of the **University of Campinas**

Acknowledgements

MeatCrossCon project that through the Strategic Research Collaboration in Food Science in the State of São Paulo, Brazil and in Denmark has financed the study with support of:

- Danish Council for Strategic Research (DCSR, process 12-1311417)
- State of São Paulo Research Foundation (FAPESP, processes 11/18228-2 and 12/50535-5)

